

- (a) 15:2
- (b) 15:1
- (e) 30:2
- (d) 30:1
- Wound caused by an injury with a blunt object, leading to tearing of soft tissue is known as
- (a) Laceration
- (b) Abrasions
- (c) Punctures
- (d) Incision
- In normal range-of-motion, movement of extremity away from midline of body is termed as
 - (a) Adduction
 - (b) Eversion
- (c) Inversion
- (d) Abduction
- 4. Which one of the following breathing patterns indicates rhythm abnormality?
 - (a) Dyspnea
 - (b) Cheye-Stokes breathing
- (c) Bradypnea
- (d) Hyperventilation

RTJL-B-NRSG - A


- 5. Which one of the following levels of illness prevention includes actions and measures that inhibit the emergence of risk factors?
 - (a) Secondary prevention
 - (b) Primordial prevention
 - (c) Primary prevention
 - (d) Tertiary prevention
- Transverse depressions in nails indicating temporary disturbance of nail growth are known as
 - (a) Koilonychia
 - (b) Paronychia .
 - (c) Clubbing
 - (d) Beau's lines
- 7. Which of the following are adventitious breath sounds on auscultation?
 - 1. Crackles
 - 2. Rhonchi
 - 3. Wheeze
 - 4. Pleural friction rub

Select the correct answer using the code given below:


- (a) 1 and 2 only
- (b) 2 and 3 only
- (c) 2, 3 and 4 only
- (d) 1, 2, 3 and 4


1


























- 2. Administration of analgesic s
- 3. Chest X-ray
- 4. Air tight dressing of the site /

Which one of the following is the correct sequence of steps of Chest tube remoyal?

(3) 3, 1, 4, 2

- (b) 2/1, 4, 3
- (c) 3, 4, 1, 2
- (d) 2\4, 1, 3
- The characteristic manifestations of Nephrotic syndrome include
 - (a) Massive proteinuria, Hypertension, Hyperlipidemia, hypoalbuminemia, foamy urine
- (b) Massive proteinuria, Hypertension, hypolipidemia, hyperalbuminemia y
- (c) Peripheral edema, hypotension, hypoalbuminemia, hypolipidemia
- (d) Hyperlipidemia, hypoalbuminemia, hypotension, cola colored urine
- 50. A patient is suffering with celiac disease. Which one of the following food items he should avoid in his diet?
 - (a) Wheat
 - (b) Eggs
 - (c) Cheese
- (d) Soy products

51. A nurse caring for a patient with Type-2 Diabetes mellitus with high dose insulin was monitoring blood glucose level between 2 am to 4 am.

(a) Dawn effect

- (b) Somogyi effect
- (c) Rebound Hypoglycemia
- (d) Insulin effectiveness A
- 52. A patient with renal dysfunction develops uremia, when his GFR is
 - (a) ≤ 15 mL/min
 - (b) > 30 mL/min
 - (c) ≥ 45 mL/min
 - (d) ≥ 60 mL/min
- 53. Consider the following statements regarding preparation of multiple-choice questions:
 - 1. Stem should be clear and explicit:
 - 2. Use negatives and double negatives
 - The alternatives should be similar in length, detail and complexity;
 - 4. Avoid using distractors that are essentially the same

Which of the statement(s) given above is/are correct?

- (a) 1 only \
- (b) 1 and 2 only
- (c) 1, 3 and 1 only/_
- (d) 1, 2, 3 and 4 7

RTJL-B-NRSG - A

10



